

Veterinär-Anatomisches Institut
der Vetsuisse-Fakultät Universität Zürich

Direktor: Prof. Dr. Alois Boos

**TRPV6 and Calbindin-D9k-expression and localisation in
the bovine uterus and placenta during pregnancy**

Inaugural-Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Nele Sprekeler

Tierärztin
aus Münster, Deutschland

genehmigt auf Antrag von

Prof. Dr. Alois Boos

Zürich 2012

Content

Content	1
Summary	3
Zusammenfassung	4
Title, Authors, Correspondence	5
Abstract	6
Keywords, Background	7
Methods	9
Animals	9
Tissue sampling and Processing	10
RNA isolation and quantitative Real-Time (TaqMan) PCR	10
Immunohistochemical analyses	11
Western blot	12
Statistics	13
Results	13
Expression of TRPV6 and Calbindin-D9k mRNA in the uterus and placenta during pregnancy	13
Localisation of TRPV6 protein in the uterus and placenta during pregnancy	14
Localisation of CaBP-9k protein in the uterus and placenta during pregnancy	14
Western blot analyses of TRPV6 and CaBP-9k protein in the uterus and placenta during pregnancy	15
Expression and localisation of TRPV6 and CaBP-9k in the placenta postpartum in cows with and without retention of fetal membranes	15
Discussion	16
Conclusion	19
Competing Interests	19
Authors contributions	20
Acknowledgements	20
References	20

Figures and figure legends	22
Figure 1 – Uterine and placental TRPV6 and CaBP-9k mRNA expression	22
Figure 2 – Immunolabelling of TRPV6 in the uterine endometrium	23
Figure 3 – Immunolabelling of TRPV6 in the placentomes and adherent fetal membranes	24
Figure 4 – Immunolabelling of CaBP-9k protein in the uterine endometrium	24
Figure 5 - Immunolabelling of CaBP-9k in placentomes and adherent fetal membranes	25
Figure 6 – Western blot analyses of endometrial and placental TRPV6 protein during pregnancy	25
Figure 7 - Western blot analyses of endometrial and placental CaBP-9k protein during pregnancy	26
Figure 8 - Placental TRPV6 and CaBP-9k mRNA expression post partum in cows with and without retained fetal membranes	27
Figure 9 – Localisation of TRPV6 protein in the placentomes and fetal membranes in cows with and without retained fetal membranes	28
Figure 10 - Localisation of placental CaBP-9k protein post partum in cows with and without retained fetal membranes	28
Figure 11 – Western blot analyses of placental TRPV6 and CaBP-9k postpartum in cows with and without retained fetal membranes	39
Primer sequences	30

Vetsuisse Faculty, University of Zurich, 2012

Nele Sprekeler

Institute of Veterinary Anatomy

TRPV6 and Calbindin-D9k-expression and localisation in the bovine uterus and placenta during pregnancy

Summary

Transient receptor potential channel type 6 (TRPV6) and Calbindin-D9k (CaBP-9k) are involved in the active calcium (Ca^{2+}) transport mechanism in many tissues including placenta and uterus, suggesting a role in the establishment and maintenance of pregnancy. Moreover, TRPV6 and CaBP-9k seem to support the materno-fetal Ca^{2+} transport that is crucial for fetal Ca^{2+} homeostasis, bone growth and development. However, it is unknown if these proteins are also involved in the aetiology of pathologies associated with parturition in cows, such as retained fetal membranes (RFM). The aim of the current study was to create an expression profile of uterine and placentomal TRPV6 and CaBP-9k mRNAs and proteins during pregnancy and postpartum in cows that discharged the fetal membranes in comparison to cows that retained them.

The results of the present study demonstrate a dynamic expression of TRPV6 and CaBP-9k during pregnancy in the bovine uterine endometrium and placentomes, suggesting a functional role for these proteins in Ca^{2+} metabolism during pregnancy. The temporal and spatial expression patterns indicate that TRPV6 and CaBP-9k may be involved in materno-fetal Ca^{2+} transport, mainly through an interplacentomal transport, and that both proteins may participate in physiological processes that are crucial for fetal and placental development. However, neither TRPV6 nor CaBP-9k seem to be causative in the retention of fetal membranes.

Keywords: TRPV6, Calbindin-D9k, cow, placenta, uterus, pregnancy

TRPV6 and Calbindin-D9k-expression and localisation in the bovine uterus and placenta during pregnancy

Zusammenfassung

Transient receptor potential channel type 6 (TRPV6) und Calbindin-D9k (CaBP-9k) sind wichtige Elemente des aktiven Calcium (Ca^{2+}) Transport in vielen Geweben und scheinen an der Etablierung und Aufrechterhaltung der Trächtigkeit beteiligt zu sein. Im Uterus sind TRPV6 und CaBP-9k als wichtige Faktoren für die myometriale Kontraktilität und Sekretion von Matrixkomponenten beschrieben. In der Plazenta scheinen beide Proteine am materno-fetalen Ca^{2+} Transport beteiligt zu sein. Ob TRPV6 und CaBP-9k eine Rolle in der Entstehung von Geburtspathologien spielen ist unbekannt.

Das Ziel dieser Studie ist ein Expressionsprofil von uterinem und plazentärem TRPV6 und CaBP-9k Protein und mRNA während der Trächtigkeit des Rindes zu erstellen sowie die plazentäre Expression von Tieren mit Nachgeburtsverhaltung (NGV) der von Tieren mit physiologischer Nachgeburtsablösung gegenüberzustellen.

Die Ergebnisse dieser Studie demonstrieren eine zeitlich dynamische Expression von TRPV6 und CaBP-9k in Uterus, Plazenta und fetalen Membranen und lassen einen Einfluss beider Proteine auf den Verlauf und Erhalt der Trächtigkeit des Rindes vermuten. Die Lokalisation sowie das zeitliche Expressionsmuster deuten auf eine Rolle beider Proteine am interplazentären materno-fetalen Ca^{2+} -Transport hin. Zudem scheinen beide Proteine einen Einfluss auf die fetale und plazentäre Entwicklung zu haben. Dennoch scheinen weder TRPV6 noch CaBP-9k eine Rolle bei der Entstehung von NGV zu spielen.

Schlüsselwörter: TRPV6, Calbindin-D9k, Trächtigkeit, Rind

TRPV6 and Calbindin-D9k-expression and localisation in the bovine uterus and placenta during pregnancy

Nele Sprekeler^{1*}, Mariusz P. Kowalewski¹, and Alois Boos¹

¹Institute of Veterinary Anatomy, Vetsuisse Faculty, University of Zurich,
Winterthurerstrasse 260, 8057 Zurich, Switzerland

*Corresponding author

Email addresses:

NS: nelesprekeler@vetanat.uzh.ch

MPK: kowalewski@vetanat.uzh.ch

AB: boosa@vetanat.uzh.ch

Abstract

Background

Transient receptor potential channel type 6 (TRPV6) and Calbindin-D9k (CaBP-9k) are involved in the active calcium (Ca²⁺) transport mechanism in many tissues including placenta and uterus, suggesting a role in the establishment and maintenance of pregnancy. Moreover, TRPV6 and CaBP-9k seem to support the materno-fetal Ca²⁺ transport that is crucial for fetal Ca²⁺ homeostasis, bone growth and development. However, it is unknown if these proteins are also involved in the aetiology of pathologies associated with parturition in cows, such as retained fetal membranes (RFM). The aim of the current study was to create an expression profile of uterine and placentomal TRPV6 and CaBP-9k mRNAs and proteins during pregnancy and postpartum in cows with and without fetal membrane release.

Methods

Uteri and placentomes of 27 cows in different stages of pregnancy and placentomes of cows with and without RFM were collected. Protein and mRNA expression of TRPV6 and CaBP-9k was investigated by real-time PCR, immunohistochemistry and Western blot.

Results

In the uterine endometrium, highest TRPV6 and CaBP-9k expression was found in the last trimester of pregnancy, with a particular increase of protein in the glandular epithelium. In the placentomes, a gradual increase in TRPV6 mRNA was detectable towards parturition, while protein expression did not change significantly.

Placentomal CaBP-9k expression did not change significantly throughout pregnancy but immunohistochemistry revealed an increase in staining intensity in the maternal crypt epithelium. Immunohistochemically, stronger placental CaBP-9k signals were seen in animals with RFM compared to animals with an undisturbed fetal membrane release, while protein levels, measured by Western blot analyses did not change significantly.

Conclusion

The results of the present study demonstrate a dynamic expression of TRPV6 and CaBP-9k during pregnancy in the bovine uterine endometrium and placentomes, suggesting a functional role for these proteins in Ca^{2+} metabolism during pregnancy. The temporal and spatial expression patterns indicate that TRPV6 and CaBP-9k may be involved in materno-fetal Ca^{2+} transport, mainly through an interplacentomal transport, and that both proteins may participate in physiological processes that are crucial for fetal and placental development. However, neither TRPV6 nor CaBP-9k seem to be causative in the retention of fetal membranes.

Keywords

TRPV6, Calbindin-D9k, cow, placenta, uterus, pregnancy

Background

The presence of Ca^{2+} in female reproductive tissues is essential during pregnancy. Uterine Ca^{2+} ions are thought to be involved in the establishment and maintenance of pregnancy, possibly by controlling myometrial activity and by influencing the secretion of extra-cellular matrix components [1-3]. Furthermore, materno-fetal transport of Ca^{2+} is crucial for fetal development, e.g. bone mineralization, neuromuscular activities, and blood coagulation [4, 5]. In the placenta, Ca^{2+} functions as an important second messenger that regulates several cellular processes such as transcription, proliferation, differentiation, necrosis and apoptosis [6]. During pregnancy, the fetal requirement for Ca^{2+} increases strikingly, especially in the last trimester of gestation: the Ca^{2+} content of bovine fetuses increased from 3g/kg at d 100 postmating to 8g/kg at d 280, indicating a high level of materno-fetal Ca^{2+} transfer [7]. This Ca^{2+} must be transported against a concentration gradient from the mother to the fetus [8], thus an active transport mechanism is required. TRPV6 and CaBP-9k (also known as S100 calcium binding protein G) are crucial elements in the active transport mechanism of mammals. TRPV6 is an epithelial Ca^{2+} channel, localised in the apical membrane of epithelial cells in several tissues,

including small intestine, kidney, uterus and placenta [9], which mediates Ca^{2+} uptake into the cell. CaBP-9k is known to facilitate intracellular Ca^{2+} transport from the apical to the basolateral cell membrane and functions additionally as an intracellular buffer of Ca^{2+} in many tissues, such as intestine, bone, uterus and placenta [10]. The expression, and to a certain extent, the regulation of CaBP-9k and TRPV6 in female reproductive tissues is well documented in several mammalian species, such as rat, mouse, pig and human [5, 11-14]. Unlike in Ca^{2+} -absorbing organs like the duodenum, where the expression of these proteins is regulated mainly via vitamin-D, their expression in the female reproductive system seems to be regulated via the steroid hormones progesterone and estrogens [2, 11, 12]. Interestingly, there are distinct variations in the expression and regulation of both proteins among different mammals, which explains their species-specific, dynamic expression patterns during the estrous cycle and pregnancy [1, 2, 11].

The clinical importance of TRPV6 in female reproductive tissues has been shown using TRPV6-knockout mice [4]. These animals showed reduced fertility, the materno-fetal Ca^{2+} transport was 40% lower and fetal Ca^{2+} accumulation was reduced compared to wildtype mice. The physiological role of uterine CaBP-9k appears to be the regulation of myometrial activity, while in the placenta CaBP-9k is thought to be involved in the regulation of the materno-fetal Ca^{2+} transport [3, 12].

Taken together, several findings indicate that TRPV6 and CaBP-9k may play an important role in the establishment and maintenance of pregnancy and the materno-fetal Ca^{2+} transport in mammals. Thus an undisturbed Ca^{2+} metabolism in the reproductive tissues is crucial for fertility and fetal health.

So far, however, the expression of uterine and placentomal TRPV6 expression has not been studied during pregnancy in the cow. Moreover, it is unknown if TRPV6 or CaBP-9k are involved in the development of pathologies related to pregnancy and parturition. The expression of bovine CaBP-9k was investigated in the bovine uterus during the estrous cycle [15], where it was mainly detected during the luteal phase, and placentomal CaBP-9k was examined in the second half of pregnancy, where its expression level increased during the last trimester, correlating with the increased fetal Ca^{2+} demand [16].

This study was performed to obtain further information about the uterine and placentomal Ca^{2+} metabolism at different stages of pregnancy in the cow. Therefore,

the temporal and spatial expression patterns of TRPV6 and CaBP-9k protein and mRNA in the uterine endometrium and in the placentomes were examined. Retained fetal membranes (RFM) in cows are described as an anomaly in the normal release of the fetal membranes within 12 hours after calving [17]. The mechanism leading to RFM is still unclear. Factors such as nutritional deficiencies, inflammation of the placenta after parturition, disturbances in maturation of the fetomaternal unit and in apoptotic processes have all been implicated in the aetiology of RFM [18]. Since Ca^{2+} plays such an important role in many cellular processes concerning placental maturation, such as proliferation, differentiation, transcription or apoptosis [6], it is also a matter of interest if TRPV6 and CaBP-9k may be involved in the detachment of the placenta postpartum. Therefore, this study further investigated the placental expression of TRPV6 and CaBP-9k postpartum in animals that discharged their fetal membranes within 12 hours after parturition and animals that did not.

Methods

Animals

Uteri of 45 pregnant Holstein-Friesian cows were collected at the local abattoir and three or four at each month 2-9 of gestation were selected at random for the present study. Within 30 minutes after the animals were killed, they were eviscerated, the pregnant uteri were opened and the crown-rump lengths of the 45 fetuses were recorded to estimate fetal age as described by Boos et al. [19].

The study also examined placental specimens of five spontaneously calving postpartum cows that discharged the fetal membranes (DFM) within 12 hours after expulsion of the fetuses and of five animals that retained fetal membranes (RFM). These cattle were members of the university dairy herd or patients from the Clinic for Bovine Obstetrics and Gynaecology, University of Veterinary Medicine Hannover, Foundation.

Animal experiments were approved by the administration of the district Hannover, and performed according to the German Law for the Protection of Animals (TierschG) and the recommendations of the German Society of Laboratory Animal Science (GV-SOLAS).

Tissue sampling and Processing

At the abattoir, at least two randomly selected and comparatively large placentomes and two segments of the interplacentomal uterine wall including adherent allantochorion were excised from each animal. Macroscopically visible caruncles, the adhering allantochorion and the surrounding segment of the uterine wall were excised from the uterine horn ipsilateral to the corpus luteum in early pregnant cows. Only a single placentome each -but no uterine wall- was collected from ten cows within one hour after the expulsion of their calf. For excision of placentomes from live cows, an elongated effeminator according to Reisinger (1906), modified by Richter (1936) (Hauptner, Solingen) was used.

A centrally located cross section of approximately 10 mm in thickness was performed on selected placentomes. The sections were cut in pieces of 10x10mm and three samples each were selected for further processing. These were: (1) uterine wall oriented – concave side – tissue cube, i.e., placental tissue including maternal plate and placental stalk material; (2) allantoid sac oriented – convex side – tissue cube, i.e., placental tissue including fetal plate material, and finally (3) tissue cube collected from the periphery – equatorial – also including fetal plate and placental tissue. The specimens were oriented and mounted in Tissue-Tek II® (Miles, Elkhart, Ind., USA) on cork lamellae, snap frozen and stored at -80°C.

For immunohistochemistry, tissue samples were rinsed in 0.9% saline, fixed in 4% neutral buffered formalin for 24 hours and finally embedded in paraffin (Histowax, Leica) as previously described [19].

RNA isolation and quantitative Real-Time (TaqMan) PCR

Total RNA from bovine uterine wall and placentomes from the 2nd to 9th month of pregnancy (n=3 animals /month) and placentomes of postpartum cows with (RFM, n=5) and without retained fetal membranes (DFM, n=6) was isolated using TRIzol Reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol. RNA concentration was measured by spectrophotometry (SmartSpecTM Plus Spectrophotometer, Biorad, Reinach, Switzerland). Purity and quality of mRNA were determined by optical density (OD) measurement. The OD 260/280 ratio of all samples was greater than 1.8. Then, 200 ng of total RNA was DNase treated (DNase I recombinant, RNase free, Roche Diagnostics, Indianapolis, USA) and reverse transcribed using the Gold RNA PCR Core Kit (Gene Amp, Applied Biosystems,

Foster City, CA). The RT reaction was run for 8 minutes at 21°C, 15 minutes at 45°C and 5 minutes at 99°C.

Real Time (TaqMan) PCR was performed using the ABI 7500 Fast Real-Time PCR System (Applied Biosystems, Rotkreuz, Switzerland), as described previously [20]. In brief, samples were amplified in duplicates in a 25- μ l reaction mixture containing 12.5 μ l FastStart Universal Probe Master (Roche Diagnostics, Mannheim, Germany), 300 nM of each primer, 200 nM TaqMan probe and 5 μ l cDNA. Autoclaved water instead of RNA and the so-called RT minus control were used as negative controls. Primers and 6-carboxyfluorescein (6-FAM)- and 6-carboxytetramethyl-rhodamine (TAMRA)-labelled probes were designed using Primer Express Software (Applied Biosystems) and ordered from Eurogentec, B-4102 Seraing, Belgium. For primer and probe sequences see Table 1. The expression levels of TRPV6 and CaBP-9k were quantified by the comparative CT method, calculated relative to the expression of the reference gene GAPDH and normalised to the calibrator. The sample with the lowest level of the respective target gene transcripts was used as the calibrator. The efficiency of the PCR reactions was verified using the CT slope method according to the instructions of the manufacturers of the ABI 7500 Fast Real-Time PCR System. The specificity of the selected PCR products was confirmed by sequencing (Microsynths, Switzerland).

Immunohistochemical analysis

Paraffin-embedded tissue samples were cut at 3 μ m. Sections were dewaxed using xylol and rehydrated through serial dilutions of ethanol to water and rinsed in water. Antigen retrieval was performed by boiling the sections in 10mM citrate buffer pH 6.0 in a microwave oven (600 W, 3 x 5 minutes) and rinsing in Tris-buffered saline (TBS, buffer stock solution: 6.1 g Trizma base, 50 ml H₂O and 37 ml 1 N HCl, diluted with H₂O to 1000 ml solution, pH adjusted to 7.6; working solution: 100 ml buffer stock solution plus 900 ml 0.85% saline). Endogenous peroxidase activity was inhibited by incubating the sections in 0.3% H₂O₂ in methanol for 15 minutes. Non-specific binding was blocked with 10% normal goat serum. The tissue sections were incubated at 4°C overnight with the primary antibody. The expression of TRPV6 was detected with a rabbit polyclonal anti-TRPV6 antibody (ACC-036, Alomone Labs, Israel) at a 1:200 dilution, which reacts with the intracellular C-terminus. The

expression of CaBP-9k was detected with a rabbit polyclonal anti-CaBP-9k-antibody (Swant, Bellinzona, Switzerland) at a dilution of 1:5000, which was produced against rat recombinant Calbindin-D9k protein. As negative controls, tissue sections were incubated with an isotype specific rabbit IgG (Vector Laboratories, Burlingame, CA, USA) and with TBS instead of the primary antibody.

After rinsing in TBS for 3x10 minutes, the biotinylated secondary antibody (PowerVision Poly-HRP IHC Detection System, goat anti-Rabbit IgG) was added in dilution of 1:200 and sections were incubated for 30 minutes at 23°C. After washing in TBS for 2x10 minutes, the tissue sections were incubated for 30 minutes with StreptABComplex/HRP Duett (Dako, Glostrup, DK), and Chromogen diaminobenzidine tetrahydrochloride (Liquid DAB+ Substrate, Dako, Baar, Switzerland) was added. Before being automatically coverslipped (RCM 2000[®]; Medite) in Pertex[®] (Medite), the sections were counterstained with haematoxylin.

Western blot

The tissue samples (n=3 animals/month) were homogenised in NET-2 buffer (50 mM TRIS-HCl, pH 7.4, 300 mM NaCl, 0.05% Nonidet P-40 [21] containing 1 µl/ml protease inhibitor (Sigma-Aldrich, Buchs, Switzerland) and centrifuged (1000 g) for 10 minutes at 4°C. The clarified supernatants were removed, and the protein concentrations were determined by the Bradford method (Bradford, 1976) (Bio-Rad Laboratories, Reinach, Switzerland). Samples were diluted with NET-2 buffer and sample buffer, containing β-mercaptoethanol, and incubated for 5 minutes at 95°C. Proteins were separated on SDS-Gel (TRPV6: 10% polyacrylamide, CaBP-9k: 12% polyacrylamide) and transferred onto PVDF membranes. For the deglycosylation of TRPV6, the proteins were denatured at 100°C and incubated with N-glycosidase F (PNGase F, New England Biolabs, Frankfurt, Germany) at 37° C for 1 hour, before separation by SDS-Gel, according to the manufacturer's protocol. After blocking the membranes in 5% fat-free milk in PBS-T (Tris-buffered saline with 0.025% Tween-20) for one hour, the membranes were incubated overnight at 4°C with the primary antibodies diluted in 2.5% milk/PBS-T. The antibodies were identical to those used for immunohistochemistry, diluted 1:500 (TRPV6) and 1:5000 (CaBP-9k). As a negative control for the TRPV6 antibody, samples were preincubated with a specific control antigen (Alomone labs, Israel).

Afterwards, the blots were washed in PBS-T three times for 10 minutes and incubated with the secondary peroxidase-conjugated antibody for one hour at 23°C (anti-rabbit IgG (H+L) HRP conjugate, Promega, Dübendorf, Switzerland). Membranes were washed five times in PBS-T for 10 minutes. Protein detection was performed by chemiluminescence (Immun-Star HRP Substrate Kit, Bio-Rad) and exposed to LAS-3000 (Fujifilm). The membranes were reblotted with anti-GAPDH antibody at a 1:5000 dilution as the loading control. The optical density of the TRPV6, CaBP-9k and GAPDH bands was quantified using ImageJ software. The values are presented as the ratio of TRPV6 and CaBP-9k optical density to the corresponding GAPDH optical density.

Statistics

The data sets of real-time and Western blot were tested for normality using the Kolmogorov-Smirnov Test [22]. Small deviations from normality were observed in the dataset of CaBP-9k mRNA expression in the placenta. Therefore, these data were analyzed by a nonparametric one-way analysis of variance using the Kruskal-Wallis test, followed by Dunn's post test. The remaining datasets were analyzed using a parametric one way analyses of variance (ANOVA) followed by Dunnetts comparison test, where the second month was taken as the control group. Real-time and Western blot data sets of animals with retained fetal membranes in comparison to animals that discharged the fetal membranes were analyzed by an unpaired one-tailed t-test. For all of the statistical analyses, the software programme GraphPad Prism 5.0 (GraphPad Software Inc., San Diego, CA, USA) was used. $P < 0.05$ was considered statistically significant. Data are presented as means \pm SD.

Results

Expression of TRPV6 and Calbindin-D9k mRNA in the uterus and placenta during pregnancy

TRPV6 and CaBP-9k mRNA expression in the uterine wall and in the placentomes from the 2nd to 9th months of pregnancy was investigated by quantitative real-time (TaqMan) PCR. In the uterine wall, TRPV6 mRNA was detected throughout pregnancy with the highest levels in the 7th and 8th months ($p < 0.05$) (Fig. 1A).

Placental TRPV6 expression levels increased from the 2nd month to the highest levels in the 7th, 8th and 9th months of pregnancy ($p < 0.05$) (Fig. 1B).

Interplacentomal uterine CaBP-9k mRNA expression was detected throughout pregnancy with a significant increase in the 7th and 8th months ($p < 0.05$) (Fig. 1C).

In the placentomes CaBP-9k mRNA was detected throughout the pregnancy without any significant changes ($p > 0.05$) (Fig. 1D).

Localisation of TRPV6 protein in the uterus and placenta during pregnancy

Immunohistochemistry was performed to determine the localisation of TRPV6 in the uterine intercaruncular wall and in the placentomes during pregnancy (Fig. 2, 3). In the uterus TRPV6 protein was detected in the luminal and in the glandular epithelium and revealed the strongest labelling in the glandular epithelium in the 3rd month and from the 7th month until parturition (Fig. 2).

In the placenta, TRPV6 was primarily expressed in the inner layer of the placentomes throughout pregnancy. Considerably lower staining was found in the middle and outer layers of the labyrinth (Fig. 3 A, D). Strongest labelling was seen in the adherent fetal membranes (Fig. 3A, C). The distribution at the cellular level was diffuse (Fig. 3B).

Localisation of CaBP-9k protein in the uterus and placenta during pregnancy

The localisation of CaBP-9k in the uterine endometrium and in the placentomes was investigated by immunohistochemistry (Fig. 4, 5). In the endometrium, CaBP-9k was detected in the luminal and glandular epithelium. The strongest staining intensity was visible in the glandular epithelium of the 7th and 8th months of pregnancy (Fig. 4C, D). Furthermore the trophoblast epithelium of the adherent fetal membranes was strongly stained (Fig. 4A, B).

Placental CaBP-9k protein was mainly detected in the luminal, inner layer and in the fetal membranes (Fig. 5A, C). Staining intensity decreased towards the outer labyrinth layer. On the cellular level, CaBP-9k was expressed in the cytoplasm and nuclei of the maternal crypt epithelium and in the uni- and binucleate fetal cells. The strongest staining was found in the binucleate cells and in the maternal crypt epithelium. In the latter the staining intensity increased towards parturition (Fig. 5B, D).

Western blot analyses of TRPV6 and CaBP-9k protein in the uterus and placenta during pregnancy

In addition to the protein localisation, the expression of TRPV6 and CaBP-9k protein in the uterine wall and in the placentomes was further investigated by Western blot analyses (Fig. 6,7). In both the uterine and placental tissues, bands of TRPV6 with a molecular mass of 75kDa were detected from the 2nd to the 9th months without significant changes in expression levels (Fig. 6). In some animals a second band of approximately 90kDa was detected in the 2nd and/or 3rd months, which represents the glycosylated variation of TRPV6.

Bands of CaBP-9k protein with a molecular weight of 9kDa were detected in the uterine and placental tissues (Fig. 7). The highest ($p < 0.05$) expression levels were found in the uterine endometrium in the 7th month of pregnancy (Fig. 7B). In the placenta, no significant changes were detectable during pregnancy (Fig. 7D).

Expression and localisation of TRPV6 and CaBP-9k in the placenta post partum in cows with and without retention of fetal membranes

The placental mRNA levels of TRPV6 and CaBP-9k of animals that retained fetal membranes postpartum compared to animals that discharged fetal membranes were examined by real-time PCR. Neither the TRPV6 nor the CaBP-9k expression levels showed significant differences between the groups ($p > 0.5$) (Fig. 8).

Immunolabelling of TRPV6 and CaBP-9k protein was performed to localise these proteins in the placentomes of animals that retained and animals that discharged the fetal membranes (Fig. 9, 10). The strongest labelling of both proteins was detected in the fetal membranes in both groups. In the placentome, TRPV6 was found in the maternal part of the placentome and in the trophoblast cells of both groups (Fig. 9B, D). Strong staining of TRPV6 was found in the maternal crypt epithelium of animals that retained their fetal membranes (Fig. 9D). Similar observations were made in the distribution of CaBP-9k in the placentome. Besides the strong staining in the fetal membranes (Fig. 10A, C), strongest signals were localised in the binucleate trophoblastic cells and in the maternal crypt epithelium of cows that retained fetal membranes (Fig. 10D). The uninucleate cells were also stained, but the intensity was considerably weaker.

In the Western blot analyses, no significant differences in TRPV6 and CaBP-9k expression were detected between the groups ($p > 0.5$) (Fig. 11).

Discussion

The results of the current study show for the first time that TRPV6 is expressed in time- and cell type specific patterns during different stages of pregnancy in the uterine endometrium and placentomes in the cow, indicating a functional role for this protein in Ca^{2+} metabolism during pregnancy. Furthermore, the expression patterns of CaBP-9k as an important intracellular Ca^{2+} transporter and buffer were investigated to check for possible colocalisation and functional correlation between both factors.

In other species, such as rodents and pigs, TRPV6 seems to participate in the implantation process, in the maintenance of pregnancy, in the fetal Ca^{2+} supply and in fetal development [1, 2, 4, 11]. CaBP-9k may play a major role in the implantation process, in the regulation of myometrial activity and in the regulation of materno-fetal Ca^{2+} transport [12, 25, 26]. The expression of CaBP-9k in the bovine endometrium and placentomes has been described previously [16, 27], however, it has not been shown to correlate with TRPV6 expression nor with pathological conditions such as fetal membrane retention.

In the current study, uterine expression of both TRPV6 and CaBP-9k mRNA was highest in the 7th and 8th months of pregnancy, the time of enhanced fetal growth and skeletal mineralization [7]. The localisation of both proteins in the glandular and luminal uterine epithelium indicates an increase in Ca^{2+} transport through or demand within these epithelia, which is met by high expression of TRPV6 and CaBP-9k.

Ca^{2+} plays a key physiological role in many cellular processes, including in secretory and exocytotic processes [28], so the increased demand in the glandular and luminal epithelia may correlate with the secretory activity of the epithelial cells. The glandular and luminal epithelia are responsible, among other functions, for the production and secretion of histiotrophe, which is essential for fetal development [29] and growth.

Thus, the localisation and the expression of TRPV6 and CaBP-9k in the uterine endometrium, mainly in the last trimester which is the time of enhanced fetal growth, indicate a functional role in secretory activities and therefore in fetal nourishment and development.

This idea is further supported by the high expression of TRPV6 and CaBP-9k in the fetal membranes, indicating a high interplacentomally active Ca^{2+} transport into the fetus, which previously has been demonstrated in-vitro by Ussing-chamber studies in

sheep [30]. The colocalisation of both proteins suggests a functional correlation between TRPV6 and CaBP-9k in the bovine endometrium, which is well documented for other epithelia such as the small intestine [31].

In the present study, the results of the Western blot analyses did not show any significant time-dependent changes in the expression of either of the proteins with the exception of uterine CaBP-9k. Moreover, distinct individual variations were detected between animals at the same stages of pregnancy, as evidenced by the high standard variations (SD) of the data obtained. The immunohistochemistry results illustrate that the measured protein (and mRNA) levels are very dependent on the anatomical composition of the samples taken for the Western blot analyses and that even small changes in anatomical structure and composition can be responsible for these variations. This could also be a reason for the divergence between the measured mRNA and protein levels, particularly since the results of the immunohistochemistry demonstrate increased protein expression in the 7th and 8th months. Nevertheless, the Western blot data show a clear trend of expression patterns and provide additional useful information. Furthermore, the antibody used for the TRPV6 Western blot analyses detected two proteins of approximately 75 and 90kDa. Since TRPV6 is a glycoprotein [32], analyses of glycosylation were performed. After incubation in the presence of N-glycosidase F, only the smaller variation of 75kDa was detectable, demonstrating that the 90kDa band represents the glycosylated variation of TRPV6. Inside the placentome, TRPV6 was mainly expressed in the luminal, inner layer of the labyrinth. While a gradual increase in placental TRPV6 mRNA was found towards parturition, the protein expression levels did not change significantly throughout pregnancy. A possible role for the discrepancy between mRNA and protein levels may be post-transcriptional regulation, or the above-mentioned possibility of small divergences in anatomical structure. A similar placental increase of TRPV6 mRNA was described in rats [2], while mouse placental TRPV6 expression peaks in the middle but not at the end of pregnancy [1]. The expression patterns of TRPV6 in the bovine placenta indicate an influence on placental Ca²⁺ metabolism, but their precise function remains unclear.

The considerably higher expression of TRPV6 protein in the interplacentomal fetal membranes compared to its expression in the placentomes provides evidence that materno-fetal Ca²⁺ transport occurs mainly in the interplacentomal regions and TRPV6 seems to play a crucial role in this process in the cow. Similar results are

documented in mice, where the intra- and extra-placental yolk sac is described to be the site of materno-fetal Ca^{2+} transport in which TRPV6 seems to be a crucial element [4].

The high expression of CaBP-9k and the colocalisation with TRPV6 in the fetal membranes supports this theory. It has been demonstrated in earlier studies that CaBP-9k levels in the ruminant interplacental trophoblasts are eight to ten times higher than in the placentomal region [27], which confirms our results and supports the idea of a facilitated diffusion of Ca^{2+} through the fetal membranes to supply the enhanced fetal need for Ca^{2+} , especially in the last trimester of pregnancy.

The intraplacental distribution of CaBP-9k, mainly at the luminal side of the labyrinth, correlates with TRPV6 localisation, but on the cellular level there are distinct variations. CaBP-9k is mostly expressed in the binucleate trophoblast cells (BNC), as described previously [16] and in the maternal crypt epithelium, while the distribution of TRPV6 is more diffuse. Interestingly, the staining intensity of CaBP-9k in the maternal epithelium increases towards parturition. The distribution of CaBP-9k and its changes indicate a functional role in placentomal Ca^{2+} metabolism. As described before, Ca^{2+} is important for many cell physiological processes, which are also crucial for placental development and degradation [6]. Thus, the high expression of CaBP-9k in the binucleate trophoblast cells may reflect the intracellular need for Ca^{2+} to support the active metabolism of these cells. In the ruminant placenta, the BNCs are, among other cells, responsible for the synthesis of hormones and proteins, for migration and cell-cell-interactions [33], all processes which are Ca^{2+} dependent [34, 35].

Furthermore, the high CaBP-9k expression in the maternal crypt epithelium possibly reflects the increased Ca^{2+} requirement for cell-specific processes, such as integrin-mediated placentation, placental growth and placental maturation [22]. The important role of intracellular Ca^{2+} as a mediator of processes concerning placental maturation in the last trimester, such as degradation, apoptosis and separation, may explain the increase of CaBP-9k expression by the maternal epithelium in the last trimester of pregnancy. Based on this theory, it is possible that disturbed Ca^{2+} cell metabolism influences maturation of the placenta and apoptotic degradation of the maternal crypt epithelium, which could lead to a disturbance in release of the fetal membranes after parturition [19, 22]. Therefore, this study further examined the expression patterns of TRPV6 and CaBP-9k in animals that retained the fetal membranes compared to

animals that discharged the fetal membranes within 12 hours after parturition. In the placentomes, both proteins were mainly detected in the maternal epithelium, CaBP-9k additionally in the trophoblast cells, especially in the binucleate cells. This localisation explains the higher expression in animals that retained the fetal membranes, since the maternal epithelium is largely intact in these animals while it is strongly reduced in cows with normal fetal membrane release. Thus, the idea that a deficiency of TRPV6 or CaBP-9k expression leads to a disturbance in release of fetal membranes could not be confirmed. Expression of both proteins reflects the status of the epithelia involvement and is rather a consequence than the cause of the retention of fetal membranes.

Conclusion

The results of the present study demonstrate for the first time that TRPV6 is dynamically expressed in the bovine uterine endometrium and in the placentomes during pregnancy, indicating a functional role in Ca^{2+} metabolism during gestation. The expression of TRPV6 and CaBP-9k in the uterine glandular epithelium, in the placental maternal epithelium and the fetal chorionic membranes suggests that these proteins are involved in materno-fetal Ca^{2+} transfer mainly through interplacental transport, especially in the last trimester of pregnancy, a time of enhanced fetal growth. Their distinct expression in the uterine glandular epithelium indicates that TRPV6 and CaBP-9k participate in uterine secretory processes, which are crucial for fetal nourishment and development.

Due to their increased expression in the maternal epithelium and in the binucleate trophoblastic cells in the placentomes, TRPV6 and CaBP-9k may function as important elements in controlling placental development, maturation and degradation through the regulation of intracellular Ca^{2+} metabolism. However, neither TRPV6 nor CaBP-9k seem to be causative in the retention of fetal membranes.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

NS was involved in the design of the study, in the coordination and performance of the experiments, in the evaluation and interpretation of the data and the drafting of the manuscript. MPK was involved in the experimental part of the project and revision of the manuscript. AB was involved in the design of the study, in the sampling procedure and supervision of the project, and was involved in drafting the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The excellent technical assistance of Urs Büchler and Elisabeth Högger-Manser is acknowledged.

References

1. Lee GS, Jeung EB: **Uterine TRPV6 expression during the estrous cycle and pregnancy in a mouse model.** *Am J Physiol Endocrinol Metab* 2007, **293**(1):E132-138.
2. Lee BM, Lee GS, Jung EM, Choi KC, Jeung EB: **Uterine and placental expression of TRPV6 gene is regulated via progesterone receptor- or estrogen receptor-mediated pathways during pregnancy in rodents.** *Reprod Biol Endocrinol* 2009, **7**:49.
3. Mathieu CL, Burnett SH, Mills SE, Overpeck JG, Bruns DE, Bruns ME: **Gestational changes in calbindin-D9k in rat uterus, yolk sac, and placenta: implications for maternal-fetal calcium transport and uterine muscle function.** *Proc Natl Acad Sci U S A* 1989, **86**(9):3433-3437.
4. Suzuki Y, Kovacs CS, Takanaga H, Peng JB, Landowski CP, Hediger MA: **Calcium channel TRPV6 is involved in murine maternal-fetal calcium transport.** *J Bone Miner Res* 2008, **23**(8):1249-1256.
5. Belkacemi L, Gariépy G, Mounier C, Simoneau L, Lafond J: **Calbindin-D9k (CaBP9k) localization and levels of expression in trophoblast cells from human term placenta.** *Cell Tissue Res* 2004, **315**(1):107-117.

6. Baczyk D, Kingdom JC, Uhlen P: **Calcium signaling in placenta.** *Cell Calcium* 2011.
7. Ferrell CL, Laster DB, Prior RL: **Mineral accretion during prenatal growth of cattle.** *J Anim Sci* 1982, **54**(3):618-624.
8. Pitkin RM: **Calcium metabolism in pregnancy and the perinatal period: a review.** *Am J Obstet Gynecol* 1985, **151**(1):99-109.
9. den Dekker E, Hoenderop JG, Nilius B, Bindels RJ: **The epithelial calcium channels, TRPV5 & TRPV6: from identification towards regulation.** *Cell Calcium* 2003, **33**(5-6):497-507.
10. Wasserman RH, Fullmer CS: **Calcium transport proteins, calcium absorption, and vitamin D.** *Annu Rev Physiol* 1983, **45**:375-390.
11. Choi Y, Seo H, Kim M, Ka H: **Dynamic expression of calcium-regulatory molecules, TRPV6 and S100G, in the uterine endometrium during pregnancy in pigs.** *Biol Reprod* 2009, **81**(6):1122-1130.
12. Choi KC, Jeung EB: **Molecular mechanism of regulation of the calcium-binding protein calbindin-D9k, and its physiological role(s) in mammals: a review of current research.** *J Cell Mol Med* 2008, **12**(2):409-420.
13. An BS, Choi KC, Kang SK, Lee GS, Hong EJ, Hwang WS, Jeung EB: **Mouse calbindin-D(9k) gene expression in the uterus during late pregnancy and lactation.** *Mol Cell Endocrinol* 2003, **205**(1-2):79-88.
14. Moreau R, Daoud G, Bernatchez R, Simoneau L, Masse A, Lafond J: **Calcium uptake and calcium transporter expression by trophoblast cells from human term placenta.** *Biochim Biophys Acta* 2002, **1564**(2):325-332.
15. Inpanbutr N, Miller EK, Petroff BK, Iacopino AM: **CaBP9K levels during the luteal and follicular phases of the estrous cycle in the bovine uterus.** *Biol Reprod* 1994, **50**(3):561-571.
16. Reiswig JD, Frazer GS, Inpanbutr N: **Calbindin-D9k expression in the pregnant cow uterus and placenta.** *Histochem Cell Biol* 1995, **104**(2):169-174.
17. Fourichon C, Seegers H, Malher X: **Effect of disease on reproduction in the dairy cow: a meta-analysis.** *Theriogenology* 2000, **53**(9):1729-1759.
18. Beagley JC, Whitman KJ, Baptiste KE, Scherzer J: **Physiology and treatment of retained fetal membranes in cattle.** *J Vet Intern Med* 2010, **24**(2):261-268.
19. Boos A, Janssen V, Mülling C: **Proliferation and apoptosis in bovine placentomes during pregnancy and around induced and spontaneous parturition as well as in cows retaining the fetal membranes.** *Reproduction* 2003, **126**(4):469-480.
20. Sprekeler N, Müller T, Kowalewski MP, Liesegang A, Boos A: **Expression patterns of intestinal calcium transport factors and ex-vivo absorption of calcium in horses.** *BMC Vet Res* 2011, **7**(1):65.
21. Ishigaki Y, Li X, Serin G, Maquat LE: **Evidence for a pioneer round of mRNA translation: mRNAs subject to nonsense-mediated decay in mammalian cells are bound by CaBP-9kP80 and CaBP-9kP20.** *Cell* 2001, **106**(5):607-617.

22. Lilliefors HW: **On the Kolmogorov-Smirnov Test for Normality with Mean and Variance Unknown.** *Journal of the American Statistical Association*, **62**(318):399.
23. Luu KC, Nie GY, Salamonsen LA: **Endometrial calbindins are critical for embryo implantation: evidence from in vivo use of morpholino antisense oligonucleotides.** *Proc Natl Acad Sci U S A* 2004, **101**(21):8028-8033.
24. Salamonsen LA, Nie G, Dimitriadis E, Robb L, Findlay JK: **Genes involved in implantation.** *Reprod Fertil Dev* 2001, **13**(1):41-49.
25. Wooding FB, Morgan G, Jones GV, Care AD: **Calcium transport and the localisation of calbindin-D9k in the ruminant placenta during the second half of pregnancy.** *Cell Tissue Res* 1996, **285**(3):477-489.
26. Berridge MJ, Bootman MD, Roderick HL: **Calcium signalling: dynamics, homeostasis and remodelling.** *Nat Rev Mol Cell Biol* 2003, **4**(7):517-529.
27. Roberts RM, Bazer FW: **The functions of uterine secretions.** *J Reprod Fertil* 1988, **82**(2):875-892.
28. Abbas: **Active transport of calcium ions across the ovine chorion in vitro.** *J Physiol (Lond)* 1993, **459**:310.
29. Hoenderop JG, Nilius B, Bindels RJ: **Calcium absorption across epithelia.** *Physiol Rev* 2005, **85**(1):373-422.
30. Hirnet D, Olausson J, Fecher-Trost C, Bodding M, Nastainczyk W, Wissenbach U, Flockerzi V, Freichel M: **The TRPV6 gene, cDNA and protein.** *Cell Calcium* 2003, **33**(5-6):509-518.
31. Wooding FB, Morgan G, Adam CL: **Structure and function in the ruminant synepitheliochorial placenta: central role of the trophoblast binucleate cell in deer.** *Microsc Res Tech* 1997, **38**(1-2):88-99.
32. Clapham DE: **Calcium signaling.** *Cell* 2007, **131**(6):1047-1058.
33. Leitinger B, McDowall A, Stanley P, Hogg N: **The regulation of integrin function by Ca(2+).** *Biochim Biophys Acta* 2000, **1498**(2-3):91-98.
34. McNaughton AP, Murray RD: **Structure and function of the bovine fetomaternal unit in relation to the causes of retained fetal membranes.** *Vet Rec* 2009, **165**(21):615-622.

Figures

Figure 1 – Uterine and placentomal TRPV6 and CaBP-9k mRNA expression

TRPV6 and CaBP-9k mRNA expression in the intercaruncular wall (Fig. 1A, C) and in the placenta ($n \geq 3$ animals/month) (Fig. 1B, D) from the 2nd to 9th months of pregnancy, determined by real-time PCR. Bars with ** differ significantly ($p < 0.05$) compared to the second month (*). Data were analysed using a parametric one-way analysis of variance (ANOVA), followed by Dunnett's comparison test and are

presented as means \pm SD. The nonparametric dataset of placental CaBP-9k was analysed using the Kruskal-Wallis test, followed by Dunn's post test.




Figure 2 – Immunolabelling of TRPV6 in the uterine endometrium

Localisation of uterine TRPV6 during pregnancy with strong labelling in the glandular (GE) and luminal epithelia (LE). Fig. 2F shows a negative control, without anti-TRPV6 treatment. Scale bars indicate 200 μ m.




Figure 3 – Immunolabelling of TRPV6 in the placentomes and adherent fetal membranes

Localisation of TRPV6 during pregnancy in the placenta and adherent membranes with strong staining in the inner layer (IL) of the placentome (A, D) and the adherent fetal membranes (FM) (A, C). Considerably weaker staining in the middle (ML) and outer labyrinth (OL) layers (A, B). Scale bars indicate 200µm.




Figure 4 – Immunolabelling of CaBP-9k protein in the uterine endometrium

Localisation of CaBP-9k in the glandular (GE) and luminal epithelia (LE) during pregnancy (A-D). Strongest staining was detected in the fetal membranes (FM) (A, B). Scale bars indicate 200µm.




Figure 5 - Immunolabelling of CaBP-9k in placentomes and adherent fetal membranes

Localisation of CaBP-9k during pregnancy with strong signals in the inner layer (IL) of the placentome and in the fetal membranes (FM) compared to the outer labyrinth layer (OL) (A, C). Strong staining in the binucleate trophoblast cells (BNC) and in the maternal epithelium (ME) inside the placentome (B, C). Fig. 2F shows a negative control, without anti-TRPV6 treatment. Scale bars indicate 200µm.




Figure 6 – Western blot analyses of endometrial and placentomal TRPV6 protein during pregnancy

A) Representative Western blot of TRPV6 protein expression in the uterine wall (n=3 animals/month). Bands of approximately 75 kDa were detected. An additional band of approximately 90 kDa was seen in the 2nd and 3rd months. After incubation with N-glycosidase F (PNGase F), only the smaller band of 75kDa was seen, demonstrating that the other band is the glycosylated variant of TRPV6. As a negative control, samples were preincubated with a control antigen.

B) Relative expression of TRPV6 to GAPDH was determined by densitometry using ImageJ software. No significant changes were detected ($p>0.05$).

C) Representative Western blot analyses of TRPV6 in the placenta from 2nd to 9th months of pregnancy ($n=3$ animals/month).

D) The expression of TRPV6 relative to GAPDH was determined by measuring the optical density using ImageJ software. No significant changes ($p>0.05$) were detected. Data were analysed using a parametric one-way analysis of variance (ANOVA), followed by Dunnett's comparison test, where the second month was taken as the control group, and are presented as means \pm SD.




Figure 7 - Western blot analyses of endometrial and placental CaBP-9k protein during pregnancy

A) Representative Western blot of CaBP-9k in the uterine wall ($n=3$ animals/month). Bands of approximately 9 kDa were detected.

B) The highest expression relative to GAPDH was found in the 7th month of pregnancy, determined by measuring the optical density using ImageJ software. Bars with ** differ significantly ($p<0.05$) compared to the 2nd month (*).

C) Representative Western blot of CaBP-9k in the placenta ($n=3$ animals/month).

D) Relative expression levels of CaBP-9k to GAPDH were determined by densitometry using ImageJ software. Significant changes in the expression levels were not detected ($p>0.05$).

Data were analysed using a parametric one-way analysis of variance (ANOVA), followed by Dunnett's comparison test, where the second month was taken as the control group, and are presented as means \pm SD.




Figure 8 - Placental TRPV6 and CaBP-9k mRNA expression post partum in cows with and without retained fetal membranes

Detection of TRPV6 (A) and Calbindin-D9k (CaBP-9k) (B) mRNA levels in the placenta of postpartum cows that retained the fetal membranes for more than 12 hours (RFM; n=5) and cows that discharged the fetal membranes (DFM; n=6), measured by real-time PCR (TaqMan). No significant changes were observed ($p>0.05$). Data were analyzed by an unpaired one-tailed t-test and are presented as means \pm SD.




Figure 9 – Localisation of TRPV6 protein in the placentomes and fetal membranes in cows with (RFM) (C, D) and without retained fetal membranes (DFM) (A, B)

Strongest labeling of TRPV6 protein was seen in the fetal membranes (FM) of all animals and in the maternal crypt epithelium (ME) of cows with RFM (D).




Figure 10 - Localisation of placental CaBP-9k protein post partum in cows with (RFM) (C, D) and without retained fetal membranes (DFM) (A, B)

Strong staining was detected in the fetal membranes (FM) of all investigated animals and in the maternal crypt epithelium (ME) of animals with retained fetal membranes (D) and in the binucleate trophoblast cells (BNC) of all animals.




Figure 11 – Western blot analyses of placental TRPV6 and CaBP-9k postpartumin cows with (RFM) and without retained fetal membranes (DFM)


A) Representative Western blot of TRPV6 in the placenta. Bands of approximately 75 kDa were detected in all animals.

B) Relative expression of TRPV6 was determined by measuring the optical density relative to GAPDH using ImageJ software. Significant changes in expression levels were not detected ($p>0.05$).

C) Representative Western blot of CaBP-9k.

D) The relative expression levels to GAPDH were determined by densitometry using ImageJ software. No significant changes were detected ($p>0.05$).

Data were analyzed by an unpaired one-tailed t-test and are presented as means \pm SD.



Tables

Table 1 – Sequences of primers and probes

Primer	Primer Sequence	Accession #
TRPV6 forward	5'-CAAGGAGCCCATGACATCTGA-3'	NM_001206189.1
TRPV6 reverse	5'-CAGGGCTTTCACGAGGTTCA-3'	
TRPV6 probe	5'-CGGCCCTGCACATAGCAGTCATGAA-3'	
CaBP-9k forward	5'-TCTCCAGAAGAAGTGAAGGGC -3'	NM_174257.2
CaBP-9k reverse	5'-CCAACACCTGGAATTCTTCG-3'	
CaBP-9k probe	5'-TCCAAGCACCCCTCGATGAGCTTTTTGA-3'	
GAPDH forward	5'-GCGATACTCACTCTTCTACCTTCGA-3'	NM_001034034.1
GAPDH reverse	5'-TCGTACCAGGAAATGAGC TTGAC-3'	
GAPDH probe	5'-CTGGCATTGCCCTCAACGACCACTT-3'	

Danksagung

In der Zeit während meiner Dissertation habe ich von vielen Menschen Unterstützung in vielerlei Hinsicht erhalten, wofür ich mich an dieser Stelle bedanken möchte.

Meinem Doktorvater, Prof. Dr. Alois Boos danke ich, dass er mir meine Dissertation in seinem Institut ermöglicht hat, mir mit seinem fachlichen Rat stets beiseite stand, für sein Verständnis und seine Unterstützung in allen Lebenslagen.

PD Dr. Mariusz P. Kowalewski für seine Einarbeitung und seine fachliche Beratung.

Urs Büchler und Elisabeth Hoegger für ihre hervorragende fachliche Hilfe bei allen Laborfragen und ihre Geduld.

Prof. Dr. Annette Liesegang für die gute Zusammenarbeit bei dem vorhergehenden Projekt und der Unterstützung bei der letzten Publikation.

Dr. Tobias Müller für die gute Zusammenarbeit und die schöne gemeinsame Zeit in der Anatomie.

Meinem lieben Team in der Anatomie, insbesondere dem Doktorandenteam Alexandra Trachsel, Aykut Gram, Barbara Fox, Corinne Schroeter, Jessica von Hof und Kathrin Nechanitzky für die vielen, abwechslungsreichen Stunden, den guten Zusammenhalt, der gegenseitigen Hilfe und Unterstützung und der Toleranz gegenüber meiner guten alten Freundin Clara.

Meinen Eltern und Rolf, dass Sie immer für mich da sind.

Meinem lieben Mann Sergey für seine moralische, fachliche und private Unterstützung und für das, was wir gemeinsam haben

Und meiner kleinen Tochter Runa, die mir meine arbeitsfreie Zeit lebendig und zuckersüß gestaltet.

Curriculum Vitae

Name, Vorname: Sprekeler, Nele
Geburtsdatum: 28.02.1978
Geburtsort: Münster, Deutschland
Nationalität: deutsch

1984-1988: Grundschule Drensteinfurt, Deutschland
1988-1996: Gymnasien Hamm und Scheessel, Deutschland
1997-1999: Fachhochschule für Kunst und Gestaltung Dr. Buhmann,
Hannover, Deutschland, Abschluss Fachabitur
1999-2001: Volkshochschule Hannover, Abendgymnasium, Abschluss
Abitur

2001-2003: Studium der Veterinärmedizin an der St. Istvan Universität,
Budapest, Ungarn
2003-2007: Studium der Veterinärmedizin an der Freien Universität Berlin,
Deutschland
Mai 2007: Approbation als Tierärztin

2009-2012: Anfertigung der Dissertation am Veterinär-Anatomischen
Institut der Vetsuisse-Fakultät Universität Zürich unter Leitung
von Prof. Dr. Alois Boos

07/2007-01/2009: Assistenztierärztin in Fachtierarztpraxis Dr. Preiss, Konstanz,
Deutschland
04/2009-07/2007: Wissenschaftliche Mitarbeiterin am Friedrich-Loeffler-Institut
für Viruskrankheiten in Tübingen, Deutschland

Zürich, den 26.07.2012